

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Hiddenwood, Walworth County
WMC-Lake-1312-000
2017

Lake Information

Name:	Hiddenwood	Maximum Depth:	17 Feet
County:	Walworth	Mean Depth:	8 Feet
Legal Description:	T124-R76-S23		
Surface Area:	27 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std frame net	June 13, 2017	5 net-nights
AFS std frame net	June 14, 2017	5 net-nights
boat shocker (night)	September 06, 2017	3600 seconds

Common Fish Species Present

Yellow Perch

Largemouth Bass

Black Crappie

Black Bullhead

Northern Pike

White Sucker

Rock Bass

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std frame net	Black Bullhead	248.0	48.1	5	1	0		85	1
	Black Crappie	139.1	43.5	2	1	2	1	92	1
	Northern Pike	0.4	0.2	100		25		84	2
	Rock Bass	0.1	0.1	100		0		123	
	White Sucker	0.1	0.1	100		0		87	
	Yellow Perch	3.2	1.7	16	10	0		96	2
boat shocker (night)	Largemouth Bass	25.0	7.0	64	15	60	15	116	3

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	
AFS std frame net	Black Bullhead										248.0	248.0
	Black Crappie										139.1	139.1
	Northern Pike										0.4	0.4
	Rock Bass										0.1	0.1
	White Sucker										0.1	0.1
	Yellow Perch										3.2	3.2
boat shocker (night)	Largemouth Bass			13.2		100.5		51.6			25.0	47.6
frame net (std 3/4 in)	Black Bullhead			4.9		8.3		7.9				7.0
	Black Crappie			0.1		2.4		1.6				1.4
	Common Carp			0.0								0.0
	Northern Pike			0.3		0.2		0.1				0.2
	White Sucker			4.5		9.9		2.3				5.6
	Yellow Perch			10.0		8.0		7.0				8.3

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017		
AFS std frame net	Black Crappie	PSD											2	
		PSD-P											2	
		Wr											92	
	Northern Pike	PSD												100
		PSD-P												25
		Wr												84
	Yellow Perch	PSD												16
		PSD-P												0
		Wr												96
frame net (std 3/4 in)	Black Crappie	PSD			100		25		100					
		PSD-P			100		0		100					
		Wr			98		107		86					
	Northern Pike	PSD			100		50		0					
		PSD-P			0		0		0					
		Wr			102		107		101					
	Yellow Perch	PSD			23		0		24					
		PSD-P			6		0		11					
		Wr			87		91		99					

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

Year Class	Age	N	Mean back-calculated length (SE) at age										
			1	2	3	4	5	6	7	8	9	10	
2014	3	19	73 (1.3)	118 (1.5)	149 (2.4)								
2008	9	2	111 (10.8)	206 (18.7)	253 (1.9)	297 (3.9)	314 (.6)	324 (1.2)	334 (3.1)	341 (1.9)	346 (2.4)		
Weighted Mean		21	77	126	159	297	314	324	334	341	346		
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20	
2014	3	19											
2008	9	2											
Weighted Mean		21											

Species: Yellow Perch

Year Class	Age	N	Mean back-calculated length (SE) at age									
			1	2	3	4	5	6	7	8	9	10
2016	1	1	97									
2014	3	6	99 (2.3)	130 (3.9)	152 (5.8)							
2013	4	15	100 (1.9)	152 (3.8)	174 (3.3)	189 (3.4)						
Weighted Mean		22	100	146	168	189						
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20
2016	1	1										
2014	3	6										
2013	4	15										
Weighted Mean		22										

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

		Mean Length (expanded sample number) at capture by age									
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	1391			156 (1363)						348 (28)	
2012	66	129 (56)		224 (10)							

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2014	0		0		16	89 (2.5)	16	84 (0.6)
	2017	1363	92 (0.7)	0		0		28	76 (1.3)


Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie
Gear: AFS std frame net



Species: Black Crappie
Gear: frame net (std 3/4 in)




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie

Gear: Frame Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2012	Largemouth Bass	Juvenile	100
2013	Black Crappie	Adult	45
2013	Largemouth Bass	Fingerling	995
2014	Largemouth Bass	Large Fingerling	680
2014	White Crappie	Adult	201
2014	Yellow Perch	Adult	250