SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Burke, Gregory County FTR-Lake-3197-000 2016

Lake Information

Name: Burke Maximum Depth: 16 Feet

County: Gregory Mean Depth: 9 Feet

Legal Description: T97-R71-S32

Surface Area: 29 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	September 29, 2016	3600 seconds
boat shocker (night)	October 05, 2016	3600 seconds

Common Fish Species Present

Largemouth Bass

Bluegill

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Quality		Preferred		Memorable		Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	Stock		Qu	Quality P		Preferred		Memorable		Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	
Blue Catfish	12	30	20	51	30	76	35	89	45	114	
Bluegill	3	8	6	15	8	20	10	25	12	30	
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30	
Brown Bullhead	5	13	8	20	11	28	14	36	17	43	
Burbot	8	20	15	38	21	53	26	67	32	82	
Channel Catfish	11	28	16	41	24	61	28	71	36	91	
Common Carp	11	28	16	41	21	53	26	66	33	84	
Flathead Catfish	14	35	20	51	28	71	34	86	40	102	
Freshwater Drum	8	20	12	30	15	38	20	51	25	63	
Gizzard Shad	7	18	11	28							
Green Sunfish	3	8	6	15	8	20	10	25	12	30	
Lake Herring	5	13	8	20	11	28	14	35	17	43	
Largemouth Bass	8	20	12	30	15	38	20	51	25	63	
Longnose Gar	16	41	27	69	36	91	45	114	55	140	
Muskellunge	20	51	30	76	38	97	42	107	50	127	
Northern Pike	14	35	21	53	28	71	34	86	44	112	
Paddlefish	16	41	26	66	33	84	41	104	51	130	
Pumpkinseed	3	8	6	15	8	20	10	25	12	30	
Redear Sunfish	4	10	7	18	9	23	11	28	13	33	
River Carpsucker	7	18	11	28	14	36	18	46	22	56	
Rock Bass	4	10	7	18	9	23	11	28	13	33	
Rudd	6	15	10	25	12	30	15	38	19	48	
Sauger	8	20	12	30	15	38	20	51	25	63	
Saugeye	9	23	14	35	18	46	22	56	27	69	
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51	
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51	
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94	
Spotted Bass	7	18	11	28	14	35	17	43	20	51	
Striped Bass	12	30	20	51	30	76	35	89	45	114	
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63	
Walleye	10	25	15	38	20	51	25	63	30	76	
White Bass	6	15	9	23	12	30	15	38	18	46	
White Crappie	5	13	8	20	10	25	12	30	15	38	
White Perch	5	13	8	20	10	25	12	30	15	38	
White Sucker	6	15	10	25	13	33	16	41	20	51	
Yellow Bass	4	10	7	18	9	23	11	28	13	33	
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36	
Yellow Perch	5	13	8	20	10	25	12	30	15	38	

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abun	dance	Stock Density Indices					Condition		
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80		
boat shocker (night)	Largemouth Bass	35.5	7.6	34	8	30		8 106	5 1		

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

		CPUE											
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg	
boat shocker (night)	Largemouth Bass		37.2			52.5			18.0		35.5	35.8	
frame net (std	Black Bullhead		10.1			12.4			8.0			7.8	
3/4 in)	Black Crappie		8.6			10.4			3.9			7.6	
	Bluegill		10.0			12.6			6.9			9.8	
	Green Sunfish					1.4			0.1			8.0	
	Northern Pike		0.4			0.4			0.2			0.3	
	Yellow Perch		0.9			0.3			2.0			1.1	

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year											
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
frame net (std	Black Crappie	PSD		45			64			72			
3/4 in)		PSD-P		0			0			8			
		Wr		102			107			104			
	Northern Pike	PSD		100			100			100			
		PSD-P		25			33			0			
		Wr		80			76			95			
	Yellow Perch	PSD		11			100			60			
		PSD-P		0			0			0			
		Wr		102			94			103			

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age												
Year	N	1	2	3	4	5	6	7	8	9	10+	
2014	75	110 (39)	165 (1)	202 (23)	239 (6)	249 (5)	243 (1)					
2011	82		153 (7)	194 (37)	217 (24)	221 (11)	233 (2)					
2008	100	121 (16)	173 (27)	197 (25)	208 (32)							

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	os		
		S-Q			Q-P	P-M		М	
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2014	11	110 (2.8)	25	102 (1.1)	3	99 (1.2)	0	

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)



Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie Gear: Frame Net

