## SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Newell City Pond, Butte County LBF-Lake-479-000 2016

## **Lake Information**

Name: Newell City Pond

County: Butte

Surface Area: 34 Acres

## **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
boat shocker (night)	June 02, 2016	2280 seconds	
frame net (std 3/4 in)	June 21, 2016	8 net-nights	
std exp gill net	June 21, 2016	1 net-nights	

# **Common Fish Species Present**

Yellow Perch

Northern Pike

Largemouth Bass

Black Crappie

Bluegill

**Channel Catfish** 

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abun	dance	St	ock De	nsity Indic	es	Coi	ndition
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	32.3	14.1	45	18	20		103	3
frame net (std 3/4 in)	Black Crappie	33.4	8.4	1		1		95	1
	Bluegill	78.6	39.6	8	1	0		101	1
	Channel Catfish	0.3	0.2	100		0		97	
	Northern Pike	1.6	0.7	92		15		84	2
	Yellow Perch	6.9	1.9	0		0		82	1
std exp gill net	Black Crappie	13.0		0		0		91	2
	Channel Catfish	2.0		100		50		94	9
	Northern Pike	3.0		100		67		90	7
	Yellow Perch	5.0		0		0		87	10

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
boat shocker (night)	Largemouth Bass							15.0			32.3	23.7
frame net (std 3/4 in)	Black Crappie							65.0			33.4	49.2
	Bluegill							35.3			78.6	57.0
	Channel Catfish										0.3	0.3
	Golden Shiner							0.0				0.0
	Largemouth Bass							0.3				0.3
	Northern Pike							0.3			1.6	1.0
	Yellow Perch							16.5			6.9	11.7
std exp gill net	Black Crappie										13.0	13.0
	Channel Catfish										2.0	2.0
	Northern Pike										3.0	3.0
	Yellow Perch										5.0	5.0
std exp gill net	Black Crappie							1.0				1.0
(150 ft)	Northern Pike							4.0				4.0
	Yellow Perch							1.0				1.0

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year										
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
frame net (std	Black Crappie	PSD							1			1
3/4 in)		PSD-P							0			1
		Wr							89			95
	Northern Pike	PSD							100			92
		PSD-P							0			15
		Wr							88			84
	Yellow Perch	PSD							0			0
		PSD-P							0			0
		Wr							84			82
std exp gill net	Black Crappie	PSD										0
		PSD-P										0
		Wr										91
	Northern Pike	PSD										100
		PSD-P										67
		Wr										90
	Yellow Perch	PSD										0
		PSD-P										0
		Wr										87
std exp gill net	Black Crappie	PSD							0			
(150 ft)		PSD-P							0			
		Wr							86			
	Northern Pike	PSD							100			
		PSD-P							0			
		Wr							82			
	Yellow Perch	PSD							0			
		PSD-P							0			
		Wr							81			

## **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	267			140 (17)	146 (8)	159 (42)	159 (200)	269 (1)			

## **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


					Length	Group	os		
			S-Q		Q-P		P-M	М	
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2013	257	90 (0.9)	3	71	0		0	
	2016	265	95 (0.7)	0		2	97 (5.3)	0	
Northern Pike Gill Net	2013	0		4	82 (2.4)	0		0	
	2016	0		1	83	1	86	1	101
Yellow Perch	2013	1	81	0		0		0	
Gill Net	2016	5	87 (7.5)	0		0		0	

#### **Length Frequency Distribution**


Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)







Species: Northern Pike Gear: std exp gill net




Species: Northern Pike Gear: std exp gill net (150 ft)



Species: Yellow Perch Gear: std exp gill net




Species: Yellow Perch Gear: std exp gill net (150 ft)




## **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.


Species: Black Crappie Gear: Frame Net



Species: Northern Pike Gear: Gill Net



Species: Yellow Perch Gear: Gill Net



# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2006	Largemouth Bass	Adult	200
2009	Largemouth Bass	Juvenile	300
2012	Largemouth Bass	Adult	100
2014	Channel Catfish	Adult	120
2014	Largemouth Bass	Juvenile	250
2014	Yellow Perch	Adult	150
2015	Rainbow Trout (Ennis)	Catchable 11"	400
2016	Rainbow Trout (Shasta)	Catchable 11"	321