

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Island South, Minnehaha County
LBS-Lake-213-001
2016

Lake Information

Name: Island South
County: Minnehaha
Surface Area: 125 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
std exp gill net	June 29, 2016	3 net-nights

Common Fish Species Present

Black Bullhead

Yellow Perch

Walleye

Common Carp

Northern Pike

Black Crappie

White Sucker

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
std exp gill net	Black Bullhead	58.0	4.7	86	4	0			
	Black Crappie	1.0	1.1	33		0	93	4	
	Common Carp	2.7	0.6	75		13			
	Northern Pike	1.3	1.7	100		50	80	2	
	Walleye	4.7	1.7	57	22	36	22	83	2
	White Sucker	0.3	0.6	100		100			
	Yellow Perch	7.7	3.3	22	14	0	88	2	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg	
		2007	2008	2009	2010	2011	2012	2013	2014	2015	2016		
large frame net	Black Bullhead			46.4		142.0		205.8					131.4
	Black Crappie			0.2		0.6		0.2					0.3
	Bluegill			1.2				0.4					0.8
	Channel Catfish			0.2									0.2
	Common Carp							0.6					0.6
	Golden Shiner			0.0									0.0
	Green Sunfish			3.2									3.2
	Northern Pike							1.4					1.4
	Sunfish Hybrid							0.0					0.0
	Walleye			0.6		0.2		0.8					0.5
	Yellow Perch			4.4		1.0		0.6					2.0
std exp gill net	Black Bullhead			19.7		115.3		156.7	151.7	101.3	58.0		100.5
	Black Crappie							1.7	0.0	0.0	1.0		0.7
	Channel Catfish			0.7									0.7
	Common Carp								1.7	5.3	2.7		3.2
	Golden Shiner								0.0	0.0			0.0
	Northern Pike			0.3		1.0		1.3	7.3	9.0	1.3		3.4
	Walleye			8.3		6.7		2.0	0.7	2.3	4.7		4.1
	White Sucker			0.3				0.3				0.3	0.3
	Yellow Perch			21.7		41.7		6.0	0.3	6.0	7.7		13.9

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
large frame net	Black Crappie	PSD			100		33		100			
		PSD-P			100		33		100			
		Wr			114		104		117			
	Northern Pike	PSD								57		
		PSD-P								0		
		Wr								103		
	Walleye	PSD			100		100		100			
		PSD-P			33		0		100			
		Wr			103		90		101			
	Yellow Perch	PSD			23		100		67			
		PSD-P			14		0		0			
		Wr			102		99		74			
std exp gill net	Black Crappie	PSD							0	0	0	33
		PSD-P							0	0	0	0
		Wr							127			93
	Northern Pike	PSD			100		33		75	91	100	100
		PSD-P			0		0		25	5	15	50
		Wr			88		97		87	85	89	80
	Walleye	PSD			96		100		83	100	100	57
		PSD-P			24		15		33	50	86	36
		Wr			97		89		92	86	96	83
	Yellow Perch	PSD			62		67		6	0	0	22
		PSD-P			22		1		6	0	0	0
		Wr			90		105		106	92	92	88

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2013	0		0		1	117	0	
Northern Pike Gill Net	2013	1	91	2	89 (3.3)	1	81	0	
	2014	2	84 (3.8)	19	85 (2.4)	1	90	0	
	2015	0		23	88 (1.3)	4	91 (2.5)	0	
	2016	0		2	79 (2.7)	2	81 (1.9)	0	
Walleye Gill Net	2013	1	94	3	95 (5.1)	2	88 (0.2)	0	
	2014	0		1	83	1	89	0	
	2015	0		1	101	5	96 (2.0)	1	87
	2016	6	82 (2.4)	3	87 (4.5)	3	82 (1.9)	2	79 (3.8)
Yellow Perch Gill Net	2013	17	107 (2.6)	0		1	92	0	
	2014	1	92	0		0		0	
	2015	18	92 (2.8)	0		0		0	
	2016	18	88 (2.0)	5	88 (3.2)	0		0	




Length Frequency Distribution


Length frequency histogram of species sampled by year.

Species: Black Crappie
Gear: large frame net




Species: Northern Pike
Gear: std exp gill net






2016


Species: Walleye
Gear: std exp gill net




2013



2014







2015



2016


Species: Yellow Perch
Gear: std exp gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: Frame Net




Species: Northern Pike
Gear: Gill Net



Species: Walleye
Gear: Gill Net



Species: Yellow Perch
Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2005	Northern Pike	Adult	532
2006	Channel Catfish	Adult	142
2007	Walleye	Adult	390
2007	Walleye	Juvenile	452
2007	Yellow Perch	Fingerling	200
2008	Walleye	Adult	106
2009	Walleye	Adult	743
2010	Walleye	Small Fingerling	14,200
2011	Walleye	Large Fingerling	478
2011	Yellow Perch	Small Fingerling	75,400
2013	Yellow Perch	Small Fingerling	138,250
2014	Walleye	Small Fingerling	14,300
2015	Walleye	Juvenile	666
2015	Walleye	Small Fingerling	5,082
2016	Walleye	Juvenile	703
2016	Walleye	Small Fingerling	5,670
2016	Yellow Perch	Adult	10,187