SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Henry, Bon Homme County LJA-Lake-588-000

2016

| Lake Information | | | |
|--------------------|------------------|----------------|---------|
| Name: | Henry | Maximum Depth: | 37 Feet |
| County: | Bon Homme | Mean Depth: | 14 Feet |
| Legal Description: | T96-R58-Sec.9-10 | | |
| Surface Area: | 104 Acres | | |
| | | | |

Surveys and Investigations

Survey methods used by gear type, date, and effort.

| Gear | Date | Effort |
|--------------------------|---------------|---------------|
| std frame net (3/8 inch) | June 01, 2016 | 10 net-nights |

Common Fish Species Present

Largemouth Bass Bluegill Black Crappie Yellow Perch White Sucker Black Bullhead Green Sunfish Common Carp Channel Catfish Walleye

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{number \ off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \, offish \ge quality \, length}{number \, of \, fish \ge stock \, length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

| | St | ock | Qu | ality | Pref | erred | Mem | orable | Tro | ophy |
|------------------|------|------|------|-------|------|-------|------|--------|------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Bigmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Black Bullhead | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| Black Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

7/16/2018 Page 3

| | St | ock | Qu | ality | Pref | erred | Mem | orable | Tro | ophy |
|----------------------------------|------|------|------|-------|------|-------|------|--------|------|----------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Blue Catfish | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Bluegill | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Bluegill X Gr. Sunfish Hybrid | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Brown Bullhead | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 36 | 17 | 43 |
| Burbot | 8 | 20 | 15 | 38 | 21 | 53 | 26 | 67 | 32 | 82 |
| Channel Catfish | 11 | 28 | 16 | 41 | 24 | 61 | 28 | 71 | 36 | 91 |
| Common Carp | 11 | 28 | 16 | 41 | 21 | 53 | 26 | 66 | 33 | 84 |
| Flathead Catfish | 14 | 35 | 20 | 51 | 28 | 71 | 34 | 86 | 40 | 102 |
| Freshwater Drum | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Gizzard Shad | 7 | 18 | 11 | 28 | | | | | | |
| Green Sunfish | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Lake Herring | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 35 | 17 | 43 |
| Largemouth Bass | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Longnose Gar | 16 | 41 | 27 | 69 | 36 | 91 | 45 | 114 | 55 | 140 |
| Muskellunge | 20 | 51 | 30 | 76 | 38 | 97 | 42 | 107 | 50 | 127 |
| Northern Pike | 14 | 35 | 21 | 53 | 28 | 71 | 34 | 86 | 44 | 112 |
| Paddlefish | 16 | 41 | 26 | 66 | 33 | 84 | 41 | 104 | 51 | 130 |
| Pumpkinseed | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Redear Sunfish | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| River Carpsucker | 7 | 18 | 11 | 28 | 14 | 36 | 18 | 46 | 22 | 56 |
| Rock Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Rudd | 6 | 15 | 10 | 25 | 12 | 30 | 15 | 38 | 19 | 48 |
| Sauger | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Saugeye | 9 | 23 | 14 | 35 | 18 | 46 | 22 | 56 | 27 | 69 |
| Shorthead Redhorse | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Smallmouth Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Smallmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Spotted Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Striped Bass | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Striped Bass Hybrid (wiper) | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Walleye | 10 | 25 | 15 | 38 | 20 | 51 | 25 | 63 | 30 | 76 |
| White Bass | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| White Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Sucker | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Yellow Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Yellow Bullhead | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 14 | 36 |
| Yellow Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

| | | Abur | idance | S | tock De | nsity India | ces | Co | ndition |
|--------------------------|-----------------|------|--------|-----|---------|-------------|-------|-----|----------------|
| Gear | Species | CPUE | CI-80 | PSD | CI-80 | PSD-P | CI-80 | Wr | CI-80 |
| std frame net (3/8 inch) | Black Bullhead | 0.2 | 0.2 | 50 |) | 0 | | | |
| | Black Crappie | 2.2 | 1.1 | 41 | 17 | 9 | | 97 | 3 |
| | Bluegill | 16.1 | 11.6 | 58 | 5 | 2 | | 107 | [′] 1 |
| | Channel Catfish | 0.1 | 0.1 | 100 |) | 0 | | 102 | ! |
| | Common Carp | 0.1 | 0.1 | 100 |) | 100 | | | |
| | Green Sunfish | 0.1 | 0.1 | 0 |) | 0 | | 83 | 5 |
| | Walleye | 0.0 | 0.0 | 0 |) | 0 | | | |
| | White Sucker | 0.4 | 0.3 | 75 | | 75 | | | |
| | Yellow Perch | 0.4 | 0.3 | 25 | i | 0 | | 135 | 52 |

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

| | | | | | | | CPUE | | | | | |
|-------------------------|-----------------|------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | Avg |
| boat shocker (night) | Largemouth Bass | 72.5 | | 35.3 | | | | 33.0 | 37.5 | 25.5 | | 40.8 |
| large frame net | Black Bullhead | 6.4 | | 2.9 | | 1.5 | | 0.7 | | | | 2.9 |
| | Black Crappie | 28.2 | | 30.3 | | 20.0 | | 16.1 | | | | 23.7 |
| | Bluegill | 28.8 | | 12.0 | | 54.4 | | 21.1 | | | | 29.1 |
| | Channel Catfish | 0.1 | | | | | | | | | | 0.1 |
| | Common Carp | 0.2 | | 0.2 | | 0.2 | | 0.4 | | | | 0.3 |
| | Green Sunfish | 0.4 | | 0.2 | | 0.1 | | | | | | 0.2 |
| | Largemouth Bass | 0.2 | | | | 0.1 | | 0.1 | | | | 0.1 |
| | Sunfish Hybrid | 0.0 | | 0.0 | | | | | | | | 0.0 |
| | White Sucker | 2.7 | | 6.2 | | 1.8 | | 3.5 | | | | 3.6 |
| | Yellow Perch | 0.3 | | 0.2 | | 0.1 | | | | | | 0.2 |
| std frame net | Black Bullhead | | | | | | | | 0.3 | | 0.2 | 0.3 |
| (3/8 inch) | Black Crappie | | | | | | | | 16.0 | 16.9 | 2.2 | 11.7 |
| | Bluegill | | | | | | | | 24.6 | 20.5 | 16.1 | 20.4 |
| | Channel Catfish | | | | | | | | 0.9 | 0.1 | 0.1 | 0.4 |
| | Common Carp | | | | | | | | | 0.1 | 0.1 | 0.1 |
| | Green Sunfish | | | | | | | | | | 0.1 | 0.1 |
| | Walleye | | | | | | | | | | 0.0 | 0.0 |
| | White Sucker | | | | | | | | 2.0 | 0.6 | 0.4 | 1.0 |
| | Yellow Perch | | | | | | | | 0.1 | | 0.4 | 0.3 |

<u>10-Year Size Structure and Condition Statistics by Gear and Species</u>

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

| | | | | | | | Ye | ar | | | | |
|-----------------|---------------|-------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | Index | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 |
| large frame net | Black Crappie | PSD | 27 | | 67 | | 6 | | 94 | | | |
| | | PSD-P | 4 | | 2 | | 0 | | 0 | | | |
| | | Wr | 101 | | 96 | | 96 | | 86 | | | |
| | Yellow Perch | PSD | 33 | | 50 | | 0 | | | | | |
| | | PSD-P | 33 | | 0 | | 0 | | | | | |
| | | Wr | 76 | | 85 | | 81 | | | | | |
| std frame net | Black Crappie | PSD | | | | | | | | 96 | 36 | 41 |
| (3/8 inch) | | PSD-P | | | | | | | | 0 | 0 | 9 |
| | | Wr | | | | | | | | 91 | 93 | 97 |
| | Walleye | PSD | | | | | | | | | | 0 |
| | | PSD-P | | | | | | | | | | 0 |
| | Yellow Perch | PSD | | | | | | | | 100 | | 25 |
| | | PSD-P | | | | | | | | 0 | | 0 |
| | | Wr | | | | | | | | 82 | | 135 |
| | | | | | | | | | | | | |

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

| | | | | Mean Len | gth (expar | nded sam | ple numbe | er) at capt | ure by age | 9 | |
|------|-----|--------------|--------------|--------------|--------------|-------------|------------|-------------|------------|---|-----|
| Year | N | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ |
| 2015 | 172 | 132 (12) | 191 (112) | | 226 (15) | 236 (34) | | | | | |
| 2014 | 161 | 138 (8) | | 230 (10) | 223 (143) | | | | | | |
| 2013 | 161 | | | 205 (56) | 213 (105) | | | | | | |
| 2011 | 204 | 145 (193) | 226 (9) | 244 (2) | | | | | | | |
| 2009 | 308 | 149 (89) | 196 (73) | 236 (126) | 238 (20) | | | | | | |
| 2007 | 282 | 152 (120) | 181 (115) | 224 (41) | 253 (7) | | 285 (2) | | | | |

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


| | | | | | Length | Group | S | | |
|----------------------------|------|-----|--------------|-----|-------------|-------|-------------|---|---------|
| | | | S-Q | | Q-P | | P-M | | М |
| Species | Year | Ν | Wr (SE) | Ν | Wr (SE) | Ν | Wr (SE) | Ν | Wr (SE) |
| Black Crappie Frame Net | 2013 | 9 | 93 (2.8) | 152 | 85 (0.5) | 0 | | 0 | |
| | 2014 | 7 | 99 (3.1) | 153 | 90 (0.6) | 0 | | 0 | |
| | 2015 | 109 | 96 (1.1) | 60 | 91 (0.7) | 0 | | 0 | |
| | 2016 | 13 | 104 (3.0) | 7 | 87 (1.8) | 2 | 83 (4.0) | 0 | |


Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: large frame net









Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.





Fish Stocking

Number of fish stocked by year, species, and size.

| Adult Adult Juvenile | 1,747 1,416 7,875 |
|----------------------------|--------------------------------|
| | - |
| Juvenile | 7,875 |
| | , |
| Large Fingerling | 3,300 |
| Small Fingerling | 40,820 |
| Fingerling | 11,400 |
| Adult | 5,445 |
| | Small Fingerling Fingerling |