### SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Long, Codington County UBS-Lake-143-000 2016

#### **Lake Information**

Name: Long Maximum Depth: 23 Feet

County: Codington

Surface Area: 2,969 Acres

#### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	July 26, 2016	4 net-nights	
AFS std gill net	July 27, 2016	4 net-nights	
AFS std gill net	July 28, 2016	4 net-nights	

# **Common Fish Species Present**

Walleye

Northern Pike

Yellow Perch

White Sucker

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

### **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abun	dance	St	ock Der	nsity Indic	ces	Co	ndition
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Walleye	10.5	2.3	60	6	48	6	89	1
	White Sucker	0.2	0.2	100		100		91	1
	Yellow Perch	32.8	4.8	18	3	4	1	103	0

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
AFS std gill net	Walleye										10.5	10.5
	White Sucker										0.2	0.2
	Yellow Perch										32.8	32.8
std exp gill net	Black Bullhead	0.5										0.5
	Green Sunfish				0.6							0.6
	Northern Pike							0.0				0.0
	Orangespotted Sunfish				0.0			0.0				0.0
	Sunfish Hybrid				0.0							0.0
	Walleye	34.5			2.5			32.0				23.0
	White Sucker							8.0				8.0
	Yellow Perch	20.0			51.4			65.2				45.5

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
AFS std gill net	Walleye	PSD										60
		PSD-P										48
		Wr										89
	Yellow Perch	PSD										18
		PSD-P										4
		Wr										103
std exp gill net	Northern Pike	PSD							0			
		PSD-P							0			
	Walleye	PSD	58			47			69			
		PSD-P	14			16			7			
		Wr	103			98			100			
	Yellow Perch	PSD	43			16			18			
		PSD-P	18			2			4			
		Wr	99			98			104			

## **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Year	Ν	1	2	3	4	5	6	7	8	9	10+
2016	125	288 (16)	348 (34)		493 (10)	543 (18)	552 (36)	597 (1)	569 (4)	627 (2)	652 (4)
2013	192	298 (59)	412 (22)	477 (99)	525 (4)	524 (3)	570 (2)	615 (1)			526 (1)
2010	45	327 (24)	433 (1)	445 (3)	470 (8)	509 (4)	546 (1)		535 (1)	629 (3)	
2007	68	374 (46)	454 (11)	462 (1)			545 (10)				

				Mean Len	gth (expa	nded sam	ple numbe	er) at capt	ure by age	)	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	393	161 (324)	238 (62)	265 (7)							
2013	428	146 (353)	223 (57)	277 (18)							
2010	928	183 (894)	203 (13)	296 (21)							

#### **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	s		
			S-Q		Q-P		P-M		M
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2013	0		0		0		0	
Walleye Gill Net	2013	60	100 (0.8)	118	100 (0.6)	14	99 (1.9)	0	
	2016	51	94 (1.0)	14	91 (1.7)	56	86 (0.7)	5	75 (5.6)
Yellow Perch Gill Net	2013	321	104 (0.4)	53	103 (0.7)	16	101 (2.1)	1	97
	2016	324	104 (0.4)	53	102 (0.8)	16	100 (1.8)	0	


#### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.


Species: Northern Pike Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net




Species: Yellow Perch Gear: std exp gill net




### **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.


Species: Northern Pike Gear: Gill Net



Species: Walleye Gear: Gill Net



Species: Yellow Perch Gear: Gill Net



# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Fry	1,500,000
2012	Walleye	Fry	1,100,000
2014	Walleye	Fry	1,450,000