SOUTH DAKOTA STATEWIDE FISHERIES SURVEY Rush South, Day County UBS-Lake-411-002 2016

Lake Information

Name: Ru	ish South
----------	-----------

County: Day

Surface Area: 1,800 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	June 21, 2016	6 net-nights
AFS std gill net	June 22, 2016	6 net-nights

Common Fish Species Present

Yellow Perch Walleye White Bass White Sucker Northern Pike Common Carp Black Crappie Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{number \ off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \, offish \ge quality \, length}{number \, of \, fish \ge stock \, length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	Stock Quality		Pref	erred	Mem	orable	Trophy		
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	Stock		Qu	ality	Pref	erred	Mem	orable	Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

		Abun	dance	St	ock Dei	Condition			
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	0.2	0.2	100		100		86	3
	Black Crappie	0.2	0.2	100		100		99	1
	Common Carp	0.3	0.3	75		75		109	6
	Northern Pike	0.5	0.4	83		0		79	4
	Walleye	5.3	1.1	50	9	11	6	88 88	1
	White Bass	2.7	0.8	91		88		92	4
	White Sucker	1.3	0.4	100		100		102	3
	Yellow Perch	9.3	1.9	85	5	52	7	7 109	1

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
AFS std gill net	Black Bullhead										0.2	0.2
	Black Crappie										0.2	0.2
	Common Carp										0.3	0.3
	Northern Pike										0.5	0.5
	Walleye										5.3	5.3
	White Bass										2.7	2.7
	White Sucker										1.3	1.3
	Yellow Perch										9.3	9.3
frame net (std	Black Bullhead				1.0			9.2				5.1
3/4 in)	Black Crappie				1.6			4.3				3.0
	Common Carp				1.3			0.2				0.8
	Northern Pike				0.3			0.5				0.4
	Orangespotted Sunfish				0.0							0.0
	Rock Bass				0.2			0.4				0.3
	Walleye				2.6			6.8				4.7
	White Bass				1.7			5.9				3.8
	White Sucker				0.6			0.2				0.4
	Yellow Perch				0.1			0.1				0.1
std exp gill net	Black Bullhead							0.8				0.8
	Black Crappie				0.0			0.5				0.3
	Common Carp				0.1							0.1
	Northern Pike				0.4			6.5				3.5
	Orangespotted Sunfish				0.0			0.0				0.0
	Spottail Shiner							0.0				0.0
	Walleye				1.4			14.7				8.1
	White Bass				0.1			0.5				0.3
	White Sucker				0.4			1.0				0.7
	Yellow Perch				4.1			12.3				8.2

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
AFS std gill net	Black Crappie	PSD										100
		PSD-P										100
		Wr										99
	Northern Pike	PSD										83
		PSD-P										0
		Wr										79
	Walleye	PSD										50
		PSD-P										11
		Wr										88
	Yellow Perch	PSD										85
		PSD-P										52
		Wr										109
frame net (std	Black Crappie	PSD				71			96			
3/4 in)		PSD-P				68			40			
		Wr				104			115			
	Northern Pike	PSD				83			89			
		PSD-P				0			11			
		Wr				81			75			
	Walleye	PSD				62			86			
		PSD-P				13			23			
		Wr				94			85			
	Yellow Perch	PSD				50			100			
		PSD-P				0			100			
		Wr				112			96			
std exp gill net	Black Crappie	PSD				0			67			
		PSD-P				0			67			
		Wr							111			
	Northern Pike	PSD				86			97			
		PSD-P				0			5			
		Wr				86			83			
	Walleye	PSD				56			86			

		Year											
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
std exp gill net	Walleye	PSD-P				16			15				
		Wr				96			89				
	Yellow Perch	PSD				73			32				
		PSD-P				32			23				
		Wr				111			98				

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age												
Year	Ν	1	2	3	4	5	6	7	8	9	10+	
2016	66	208 (2)	304 (27)	386 (5)	414 (14)	479 (4)	506 (10)	530 (1)	527 (1)	571 (1)	705 (1)	
2013	93	130 (5)	302 (3)	419 (55)	492 (14)	517 (12)	633 (1)		600 (1)		580 (1)	
2010	28	219 (3)	349 (10)	429 (10)	475 (1)	549 (4)						

Species: Yellow Perch

	Mean Length (expanded sample number) at capture by age												
Year	N	1	2	3	4	5	6	7	8	9	10+		
2016	114	135 (8)	208 (33)	253 (37)	287 (7)	293 (24)	333 (2)	339 (2)	340 (1)				
2013	76	102 (2)	178 (55)	263 (10)	300 (6)		323 (3)						
2010	81	127 (18)	207 (35)	255 (28)									

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


					Length	Group	S		
			S-Q		Q-P	1	P-M	М	
Species	Year	Ν	Wr (SE)	Ν	Wr (SE)	N	Wr (SE)	Ν	Wr (SE)
Black Crappie Frame Net	2013	3	128 (3.5)	41	117 (1.0)	26	113 (1.6)	3	104 (3.4)
Northern Pike Gill Net	2013	1	75	36	83 (1.1)	2	76 (4.0)	0	
	2016	1	89	5	77 (2.8)	0		0	
Walleye Gill Net	2013	12	91 (1.6)	63	89 (0.8)	12	86 (1.4)	1	87
	2016	32	88 (0.8)	25	89 (0.9)	6	87 (1.3)	1	74
Yellow Perch Gill Net	2013	50	99 (0.9)	7	102 (3.4)	11	97 (1.6)	6	92 (3.8)
	2016	17	114 (1.9)	37	111 (0.9)	46	108 (1.1)	12	100 (1.9)


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie Gear: frame net (std 3/4 in)









Species: Northern Pike Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net





Species: Yellow Perch Gear: std exp gill net




Historic Fish Sizes and Relative Abundance


Size distribution per net by color for species sampled by year.





Species: Northern Pike Gear: Gill Net





Species: Yellow Perch Gear: Gill Net

