SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Krause East, Day County UJA-Lake-866-024 2016

Lake Information

Name: Krause East Maximum Depth: 18 Feet

County: Day

Surface Area: 171 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	August 23, 2016	6 net-nights

Common Fish Species Present

Yellow Perch Walleye

Northern Pike

Black Bullhead

Common Carp

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abun	Abundance Stock Density Indices				es	Condition		
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80	
AFS std gill net	Black Bullhead	2.7	2.8	50	21	13		95	4	
	Common Carp	0.5	0.5	33		33		112	9	
	Walleye	0.7	0.3	75		50		98	4	
	Yellow Perch	19.7	3.9	0		0		92	1	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
AFS std gill net	Black Bullhead										2.7	2.7
	Common Carp										0.5	0.5
	Walleye										0.7	0.7
	Yellow Perch										19.7	19.7
frame net (std	Black Bullhead							24.6				24.6
3/4 in)	Black Crappie							0.3				0.3
	Common Carp							0.0				0.0
	Green Sunfish							1.9				1.9
	Walleye							3.3				3.3
	Yellow Perch							5.6				5.6
std exp gill net	Black Bullhead	3.7						33.7				18.7
	Black Crappie	0.7										0.7
	Common Carp							0.0				0.0
	Saugeye	0.3										0.3
	Walleye	1.3						8.0				4.7
	Yellow Perch	65.3						14.0				39.7

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Υe	ar				
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
AFS std gill net	Walleye	PSD										75
		PSD-P										50
		Wr										98
	Yellow Perch	PSD										0
		PSD-P										0
		Wr										92
frame net (std	Black Crappie	PSD							33			
3/4 in)		PSD-P							33			
		Wr							110			
	Walleye	PSD							30			
		PSD-P							5			
		Wr							92			
	Yellow Perch	PSD							3			
		PSD-P							1			
		Wr							88			
std exp gill net	Black Crappie	PSD	0									
		PSD-P	0									
		Wr	111									
	Walleye	PSD	100						25			
		PSD-P	0						0			
		Wr	87						97			
	Yellow Perch	PSD	7						17			
		PSD-P	4						0			
		Wr	101						97			
		PSD PSD-P Wr PSD PSD-P	100 0 87 7 4						0 97 17 0			

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	4	363 (1)	468 (1)				588 (2)				
2013	24	290 (18)		467 (6)							
2007	4			438 (4)							

Species: Yellow Perch

	Mean Length (expanded sample number) at capture by age										
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	118	163 (117)	194 (1)								
2013	42	153 (33)	217 (9)								

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

	,		Length Groups									
			S-Q		Q-P		P-M		М			
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)			
Black Crappie Frame Net	2013	2	117 (20.2)	0		0		1	96			
Walleye Gill Net	2013	18	95 (1.2)	6	100 (1.4)	0		0				
	2016	1	98	1	91	1	98	1	105			
Yellow Perch Gill Net	2013	35	96 (1.0)	7	102 (9.8)	0		0				
	2016	118	92 (0.7)	0		0		0				


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net




Species: Yellow Perch Gear: std exp gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie Gear: Frame Net



Species: Walleye Gear: Gill Net



Species: Yellow Perch Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2005	Walleye	Fingerling	2,280
2008	Gizzard Shad	Adult	50
2008	Walleye	Fry	200,000
2008	Yellow Perch	Fry	183,000
2009	Gizzard Shad	Adult	50
2009	Walleye	Fry	200,000
2009	Yellow Perch	Fry	188,000
2010	Black Crappie	Fingerling	10,080
2010	Walleye	Fry	200,000
2012	Walleye	Fry	90,000
2014	Walleye	Fry	100,000