# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Drywood North, Roberts County UMN-Lake-476-800 2016

### **Lake Information**

Name: Drywood North Maximum Depth: 15 Feet

County: Roberts

Surface Area: 2,825 Acres

### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	May 24, 2016	4 net-nights	
AFS std gill net	May 25, 2016	4 net-nights	
AFS std gill net	May 26, 2016	4 net-nights	

# **Common Fish Species Present**

Yellow Perch

Walleye

Northern Pike

Black Bullhead

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Quality		Preferred		Memorable		Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

# **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abur	Abundance			Stock Density Indices				
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80	
AFS std gill net	Black Bullhead	76.4	13.3	22	2	19		2 98	1	
	Northern Pike	1.4	0.5	100		24		90	3	
	Walleye	2.7	0.9	88		50	14	4 88	1	
	Yellow Perch	1.7	0.6	85		45	18	3 97	3	

# 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
AFS std gill net	Black Bullhead										76.4	76.4
	Northern Pike										1.4	1.4
	Walleye										2.7	2.7
	Yellow Perch										1.7	1.7
frame net (std	Black Bullhead				59.2			210.6				134.9
3/4 in)	Northern Pike				0.5			0.4				0.5
	Walleye				0.3			1.8				1.1
	Yellow Perch				5.3							5.3
std exp gill net	Black Bullhead	66.7	5.3		2.2			39.0				28.3
	Northern Pike	0.7	5.7		1.4			11.7				4.9
	Walleye	3.3	6.7		2.6			6.0				4.7
	Yellow Perch	42.0	14.0		4.8			2.7				15.9

# 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Υe	ear				
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
AFS std gill net	Northern Pike	PSD										100
		PSD-P										24
		Wr										90
	Walleye	PSD										88
		PSD-P										50
		Wr										88
	Yellow Perch	PSD										85
		PSD-P										45
		Wr										97
frame net (std	Northern Pike	PSD				100			100			
3/4 in)		PSD-P				56			33			
		Wr				86			80			
	Walleye	PSD				100			45			
		PSD-P				80			16			
		Wr				91			83			
	Yellow Perch	PSD				1						
		PSD-P				0						
		Wr				102						
std exp gill net	Northern Pike	PSD	100	35		60			79			
		PSD-P	100	35		16			9			
		Wr	93	97		99			82			
	Walleye	PSD	100	10		50			83			
		PSD-P	50	5		20			44			
		Wr	107	94		97			85			
	Yellow Perch	PSD	15	33		16			31			
		PSD-P	1	2		0			0			
		Wr	97	86		99			87			

# **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

				Mean Ler	ngth (expa	nded sam	ple numb	er) at capt	ture by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	33		210 (1)		352 (5)	444 (1)	470 (14)	582 (9)	632 (2)		665 (1)
2013	36	129 (1)	240 (2)	317 (5)	496 (16)	532 (8)		640 (1)		640 (1)	660 (2)
2010	54	208 (6)	347 (24)	415 (6)	510 (16)			616 (1)	600 (1)		
2008	42	222 (23)	363 (17)	473 (1)		559 (1)					
2007	23	212 (13)	400 (1)		516 (8)		565 (1)				

Species: Yellow Perch

			I	Mean Len	gth (expa	nded sam	ple numb	er) at capt	ure by age	9	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	20		164 (2)	225 (7)	236 (3)	274 (2)	301 (2)	287 (4)			
2013	16			168 (2)	194 (12)	198 (2)					
2010	347	91 (18)	113 (283)	178 (34)	180 (11)	184 (2)					

### **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

			Length Groups										
		S-Q			Q-P		P-M		М				
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)				
Northern Pike Gill Net	2013	15	80 (1.7)	49	83 (0.9)	6	86 (4.4)	0					
	2016	0		13	90 (3.0)	3	97 (6.0)	1	80				
Walleye Gill Net	2013	6	79 (3.0)	14	86 (1.2)	13	87 (1.5)	3	86 (3.4)				
	2016	4	92 (2.2)	12	85 (1.5)	14	89 (1.7)	2	87 (6.9)				
Yellow Perch Gill Net	2013	11	88 (0.6)	5	83 (4.3)	0		0					
	2016	3	107 (3.6)	8	103 (2.4)	7	91 (2.9)	2	84 (2.0)				


### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.


Species: Northern Pike Gear: AFS std gill net




Species: Northern Pike Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net




Species: Yellow Perch Gear: std exp gill net




# **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.

Species: Northern Pike Gear: Gill Net



Species: Walleye Gear: Gill Net



Species: Yellow Perch Gear: Gill Net

