SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

George, Lake County VER-Lake-81-000 2016

Lake Information

Name:	George	Maximum Depth:	9 Feet
County:	Lake	Mean Depth:	4 Feet
Legal Description:	T105-R54-Sec. 15, 16, 21		
Surface Area:	245 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
std exp gill net	August 17, 2016	3 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

Green Sunfish

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{number \ off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \, offish \ge quality \, length}{number \, of \, fish \ge stock \, length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	 ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

		Abun	Abundance			Stock Density Indices				
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80	
std exp gill net	Black Bullhead	5.3	1.3	100		50	21			
	Green Sunfish	0.3	0.6	0		0				
	Walleye	82.3	3.3	9	3	4	2	80) 1	
	Yellow Perch	17.7	8.2	89	7	55	10	100	2	

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

10-Year Catch Per Unit Effort by Gear and Species

							CPUE					
Gear	Species	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Avg
large frame net	Black Bullhead						275.4					275.4
	Green Sunfish						2.0					2.0
	Walleye						35.4					35.4
	Yellow Perch						6.0					6.0
std exp gill net	Black Bullhead					1.7	9.7		5.0	24.0	5.3	9.1
	Green Sunfish								0.3		0.3	0.3
	Walleye					25.3	85.3		40.7	11.7	82.3	49.1
	Yellow Perch					27.7	39.3		4.0	9.0	17.7	19.5

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
large frame net	Walleye	PSD						13				
		PSD-P						0				
		Wr						93				
	Yellow Perch	PSD						77				
		PSD-P						17				
		Wr						115				
std exp gill net	Walleye	PSD					4	8		81	69	9
		PSD-P					4	0		3	14	4
		Wr						94		99	82	80
	Yellow Perch	PSD					41	76		42	100	89
		PSD-P					0	20		25	37	55
		Wr						112		104	102	100

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2014	12	176 (5)	168 (2)	262 (3)	215 (2)						

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


					Length	Group	S		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Walleye Gill Net	2012	235	94 (0.5)	21	90 (1.7)	0		0	
	2014	23	105 (1.6)	95	97 (0.7)	4		0	
	2015	11	81	19	83 (3.4)	5	81 (1.9)	0	
	2016	225	82 (0.8)	11	59 (1.5)	11	73 (3.1)	0	
Yellow Perch Gill Net	2012	28	117 (1.3)	66	112 (0.8)	24	107 (1.2)	0	
	2014	7	107 (4.5)	2	99 (11.5)	3	99 (0.9)	0	
	2015	0		17	103 (1.7)	8	106 (4.6)	2	79 (4.3)
	2016	6	101	18	100 (2.7)	26	102 (2.3)	3	81 (3.1)

Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Walleye Gear: std exp gill net







Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.





Species: Yellow Perch Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2006	Walleye	Fry	1,200,000
2007	Walleye	Small Fingerling	28,060
2010	Walleye	Small Fingerling	51,200
2010	Yellow Perch	Fingerling	112,258
2011	Walleye	Small Fingerling	25,100
2011	Yellow Perch	Fry	2,065,000
2014	Walleye	Fry	233,000