

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Island South, Minnehaha County**  
**LBS-Lake-213-001**  
**2015**

**Lake Information**

**Name:** Island South  
**County:** Minnehaha  
**Surface Area:** 125 Acres

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
std exp gill net	June 24, 2015	3 net-nights

## **Common Fish Species Present**

---

Black Bullhead

Northern Pike

Yellow Perch

Common Carp

Walleye

Golden Shiner

Black Crappie

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
std exp gill net	Black Bullhead	101.3	14.9	27	3	0			
	Black Crappie	0.0	0.0	0		0			
	Common Carp	5.3	7.3	13		0			
	Golden Shiner	0.0	0.0						
	Northern Pike	9.0	3.3	100		15	89	2	
	Walleye	2.3	1.3	100		86	96	3	
	Yellow Perch	6.0	2.2	0		0	92	4	

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
large frame net	Black Bullhead				46.4		142.0		205.8			131.4
	Black Crappie				0.2		0.6		0.2			0.3
	Bluegill				1.2				0.4			0.8
	Channel Catfish				0.2							0.2
	Common Carp								0.6			0.6
	Golden Shiner				0.0							0.0
	Green Sunfish				3.2							3.2
	Northern Pike								1.4			1.4
	Sunfish Hybrid								0.0			0.0
	Walleye				0.6		0.2		0.8			0.5
	Yellow Perch				4.4		1.0		0.6			2.0
std exp gill net	Black Bullhead				19.7		115.3		156.7	151.7	101.3	108.9
	Black Crappie								1.7	0.0	0.0	0.6
	Channel Catfish				0.7							0.7
	Common Carp									1.7	5.3	3.5
	Golden Shiner									0.0	0.0	0.0
	Northern Pike				0.3		1.0		1.3	7.3	9.0	3.8
	Walleye				8.3		6.7		2.0	0.7	2.3	4.0
	White Sucker				0.3				0.3			0.3
	Yellow Perch				21.7		41.7		6.0	0.3	6.0	15.1

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2006	2007	2008	2009	2010	2011	2012	2013	2014	2015		
large frame net	Black Crappie	PSD				100		33		100				
		PSD-P				100		33		100				
		Wr				114		104		117				
	Northern Pike	PSD									57			
		PSD-P									0			
		Wr									103			
	Walleye	PSD				100		100		100				
		PSD-P				33		0		100				
		Wr				103		90		101				
	Yellow Perch	PSD				23		100		67				
		PSD-P				14		0		0				
		Wr				102		99		74				
std exp gill net	Black Crappie	PSD									0	0	0	
		PSD-P									0	0	0	
		Wr									127			
	Northern Pike	PSD				100		33		75	91	100		
		PSD-P				0		0		25	5	15		
		Wr				88		97		87	85	89		
	Walleye	PSD				96		100		83	100	100		
		PSD-P				24		15		33	50	86		
		Wr				97		89		92	86	96		
	Yellow Perch	PSD				62		67		6	0	0		
		PSD-P				22		1		6	0	0		
		Wr				90		105		106	92	92		

## **Fish Condition**

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2011	2	108 (1.4)	0		0		1	95
	2013	0		0		1	117	0	
Northern Pike Gill Net	2011	2	100 (3.1)	1	92	0		0	
	2013	1	91	2	89 (3.3)	1	81	0	
	2014	2	84 (3.8)	19	85 (2.4)	1	90	0	
	2015	0		23	88 (1.3)	4	91 (2.5)	0	
Walleye Gill Net	2011	0		17	90 (1.4)	3	87 (2.5)	0	
	2013	1	94	3	95 (5.1)	2	88 (0.2)	0	
	2014	0		1	83	1	89	0	
	2015	0		1	101	5	96 (2.0)	1	87
Yellow Perch Gill Net	2011	41	109 (1.7)	83	104 (1.2)	1	79	0	
	2013	17	107 (2.6)	0		1	92	0	
	2014	1	92	0		0		0	
	2015	18	92 (2.8)	0		0		0	





# Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Black Crappie



Gear: large frame net






Species: Northern Pike


Gear: std exp gill net









Species: Walleye  
 Gear: std exp gill net





Species: Yellow Perch  
Gear: std exp gill net







## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie  
Gear: Frame Net




Species: Northern Pike  
Gear: Gill Net



Species: Walleye  
Gear: Gill Net



Species: Yellow Perch  
Gear: Gill Net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2005	Northern Pike	Adult	532
2006	Channel Catfish	Adult	142
2007	Walleye	Adult	390
2007	Walleye	Juvenile	452
2007	Yellow Perch	Fingerling	200
2008	Walleye	Adult	106
2009	Walleye	Adult	743
2010	Walleye	Small Fingerling	14,200
2011	Walleye	Large Fingerling	478
2011	Yellow Perch	Small Fingerling	75,400
2013	Yellow Perch	Small Fingerling	138,250
2014	Walleye	Small Fingerling	14,300
2015	Walleye	Juvenile	666
2015	Walleye	Small Fingerling	5,082