SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Menno, Hutchinson County LJA-Lake-52-000 2015

Lake Information

Name: Menno Maximum Depth: 34 Feet

County: Hutchinson Mean Depth: 13 Feet

Legal Description: T98N-R57W-Sec. 32

Surface Area: 39 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	June 02, 2015	4800 seconds
std frame net (3/8 inch)	August 19, 2015	10 net-nights

Common Fish Species Present

Largemouth Bass

Bluegill

Black Crappie

Black Bullhead

White Sucker

Green Sunfish

Yellow Perch

Channel Catfish

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

		Abun	dance	St	ock Der	nsity India	ces	Co	ndition
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	48.0	9.8	64	9	64	9	94	. 1
std frame net (3/8 inch)	Black Bullhead	3.2	1.7	88		84	10)	
	Black Crappie	13.3	4.5	89	4	71	6	101	1
	Bluegill	13.7	5.9	100		91	4	104	. 1
	Channel Catfish	0.1	0.1	100		0			
	Green Sunfish	0.3	0.4	0		0		90	4
	Largemouth Bass	0.2	0.2	0		0		93	1
	White Sucker	0.9	0.8	100		100			
	Yellow Perch	0.1	0.1	100		0		95	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Avg
boat shocker (night)	Largemouth Bass		70.0		33.0		112.0		84.0	29.1	48.0	62.7
large frame net	t Black Bullhead		7.8		10.4		5.0		10.8			8.5
	Black Crappie		32.5		4.3		29.4		20.6			21.7
	Bluegill		23.3		43.1		105.8		43.0			53.8
	Channel Catfish		0.1		0.1							0.1
	Green Sunfish		0.5				0.2					0.4
	Largemouth Bass				0.2		0.2		0.1			0.2
	Sunfish Hybrid		0.0		0.0		0.0					0.0
	White Sucker		0.4		1.3		8.0		1.2			0.9
	Yellow Perch				2.6		1.9		1.9			2.1
std frame net	Black Bullhead									3.0	3.2	3.1
(3/8 inch)	Black Crappie									27.0	13.3	20.2
	Bluegill									27.2	13.7	20.5
	Channel Catfish									0.2	0.1	0.2
	Green Sunfish										0.3	0.3
	Largemouth Bass									0.0	0.2	0.1
	White Sucker									0.4	0.9	0.7
	Yellow Perch									0.4	0.1	0.3

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
large frame net	Black Crappie	PSD		75		56		48		40		
		PSD-P		1		0		3		0		
		Wr		102		111		97		87		
	Yellow Perch	PSD				77		0		58		
		PSD-P				0		0		0		
		Wr				91		89		75		
std frame net	Black Crappie	PSD									100	89
(3/8 inch)		PSD-P									18	71
		Wr									106	101
	Yellow Perch	PSD									100	100
		PSD-P									0	0
		Wr									95	95

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

					Me	an back-	calculated	length (SE) at ag	е		
Year Class	Age	N	1	2	3	4	5	6	7	8	9	10
2014	1	9	88 (3.3)									
2013	2	4	89 (3.9)	175 (11.3)								
2010	5	17	71 (1.4)	122 (2.1)	163 (1.8)	206 (2.1)	239 (2.6)					
Weighted Mean		30	79	132	163	206	239					
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20
2014	1	9										
2013	2	4										
2010	5	17										
Weighted Mean		30										

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

	,			Mean Len	gth (expa	nded sam	ple numb	er) at capt	ure by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2015	133	162 (14)	218 (6)			255 (113)					
2014	135			212 (1)	242 (134)						
2013	206	145 (2)		197 (204)							
2011	376	134 (189)	209 (129)	232 (53)	280 (6)						
2009	45	131 (6)	201 (38)	220 (1)							
2007	325	136 (20)	185 (33)	207 (207)	219 (22)		223 (6)	241 (15)	234 (19)	235 (6)	

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

			Length Groups										
			S-Q	1	Q-P		P-M		М				
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)				
Black Crappie Frame Net	2011	152	102 (0.8)	133	92 (0.4)	9	88 (3.2)	0					
	2013	123	84 (2.3)	83	88 (2.0)	0		0					
	2014	0		111	106 (0.6)	24	105 (0.4)	0					
	2015	15	110 (2.1)	24	100 (1.3)	94	98 (0.5)	0					

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: large frame net



Species: Black Crappie Gear: std frame net (3/8 inch)

11

13

15

17

19

Total Length (cm)

21


23

25

27

29

20 | 0 | 9




Total Length (cm)

Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie Gear: Frame Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2004	Channel Catfish	Adult	170
2005	Channel Catfish	Adult	100
2006	Channel Catfish	Adult	50
2006	Largemouth Bass	Adult	95
2013	Channel Catfish	Large Fingerling	4,950