

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Henry, Bon Homme County

LJA-Lake-588-000

2015

Lake Information

Name: Henry **Maximum Depth:** 37 Feet
County: Bon Homme **Mean Depth:** 14 Feet
Legal Description: T96-R58-Sec.9-10
Surface Area: 104 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	May 26, 2015	7200 seconds
std frame net (3/8 inch)	August 18, 2015	10 net-nights

Common Fish Species Present

Largemouth Bass

Bluegill

Black Crappie

White Sucker

Common Carp

Channel Catfish

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	25.5	6.3	78	9	29	9	92	2
std frame net (3/8 inch)	Black Crappie	16.9	9.7	36	5	0		93	1
	Bluegill	20.5	7.5	84	4	2		90	1
	Channel Catfish	0.1	0.1	100		0		91	
	Common Carp	0.1	0.1	100		100			
	White Sucker	0.6	0.4	100		100			

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
boat shocker (night)	Largemouth Bass		72.5		35.3				33.0	37.5	25.5	40.8
large frame net	Black Bullhead		6.4		2.9		1.5		0.7			2.9
	Black Crappie		28.2		30.3		20.0		16.1			23.7
	Bluegill		28.8		12.0		54.4		21.1			29.1
	Channel Catfish		0.1									0.1
	Common Carp		0.2		0.2		0.2		0.4			0.3
	Green Sunfish		0.4		0.2		0.1					0.2
	Largemouth Bass		0.2				0.1		0.1			0.1
	Sunfish Hybrid		0.0		0.0							0.0
	White Sucker		2.7		6.2		1.8		3.5			3.6
	Yellow Perch		0.3		0.2		0.1					0.2
std frame net (3/8 inch)	Black Bullhead									0.3		0.3
	Black Crappie									16.0	16.9	16.5
	Bluegill									24.6	20.5	22.6
	Channel Catfish									0.9	0.1	0.5
	Common Carp										0.1	0.1
	White Sucker									2.0	0.6	1.3
	Yellow Perch									0.1		0.1

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
large frame net	Black Crappie	PSD		27		67		6		94			
		PSD-P		4		2		0		0			
		Wr		101		96		96		86			
	Yellow Perch	PSD		33		50		0					
		PSD-P		33		0		0					
		Wr		76		85		81					
std frame net (3/8 inch)	Black Crappie	PSD									96	36	
		PSD-P									0	0	
		Wr									91	93	
	Yellow Perch	PSD										100	
		PSD-P										0	
		Wr										82	

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2014	1	7	69 (2.2)											
2013	2	18	84 (2.9)	149 (3.3)										
2011	4	7	90 (2.6)	156 (2.7)	201 (3.7)	216 (3)								
2010	5	12	90 (2.8)	157 (3.7)	199 (3.1)	214 (2.9)	225 (2.7)							
Weighted Mean		44	84	153	200	215	225							
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20		
2014	1	7												
2013	2	18												
2011	4	7												
2010	5	12												
Weighted Mean		44												

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Year	N	Mean Length (expanded sample number) at capture by age									
		1	2	3	4	5	6	7	8	9	10+
2015	172	132 (12)	191 (112)		226 (15)	236 (34)					
2014	161	138 (8)		230 (10)	223 (143)						
2013	161			205 (56)	213 (105)						
2011	204	145 (193)	226 (9)	244 (2)							
2009	308	149 (89)	196 (73)	236 (126)	238 (20)						
2007	282	152 (120)	181 (115)	224 (41)	253 (7)		285 (2)				

Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).



Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2011	189	97 (0.8)	11	92 (1.5)	0		0	
	2013	9	93 (2.8)	152	85 (0.5)	0		0	
	2014	7	99 (3.1)	153	90 (0.6)	0		0	
	2015	109	96 (1.1)	60	91 (0.7)	0		0	

Length Frequency Distribution



Length frequency histogram of species sampled by year.

Species: Black Crappie

Gear: large frame net




Species: Black Crappie
Gear: std frame net (3/8 inch)



Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie
Gear: Frame Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2004	Black Crappie	Adult	1,829
2004	Bluegill	Adult	1,510
2004	Channel Catfish	Adult	500
2004	Largemouth Bass	Adult	1,029
2004	Largemouth Bass	Fingerling	400
2004	Yellow Perch	Adult	1,016
2004	Yellow Perch	Fingerling	2,448
2011	Yellow Perch	Adult	1,747
2012	Yellow Perch	Adult	1,416
2012	Yellow Perch	Juvenile	7,875
2013	Channel Catfish	Large Fingerling	3,300
2014	Yellow Perch	Small Fingerling	40,820
2015	Largemouth Bass	Fingerling	11,400